

Analysis of Volatile Oil of the Stem Bark of *Cinnamomum Zeylanicum* and its antimicrobial activity

Shahnaz S. Husain, Mohammed Ali*

Phytochemistry Research Laboratory, Faculty of Pharmacy, JamiaHamdard, Hamdard Nagar,
New Delhi – 110 062, India

ABSTRACT

*Cinnamomumzeylanicum*Blume (Lauraceae)is a moderate sized tree native to Sri Lanka and southern India. Its stembark is used to treat stomach cramps, gastric irritation, dysentery, diarrhoea, nausea, neuralgia, rheumatism, toothache and paralysis of the tongue. The bark essential oil was characterized mainly by cinnamic aldehyde (91.5 %). There were thirteen monoterpenes (5.9 %) and ten of them were monoterpene hydrocarbons (5.1 %). The predominant monoterpene was *l*-limonene (3.1 %). There was three sesquiterpenes, viz. 3, 7-guaiadene, *cis*-calamenene and caryophyllene oxide, all of them were present in trace amounts. In addition to cinnamic aldehyde, the other aromatic components detected were benzaldehyde, *o*-cymene, eugenol , (E)-methyl cinnamate and benzyl benzoate, all of them occurred in lesser than 1 % yield. The oil was devoid of any aliphatic constituent. The volatile oil showed significant antimicrobial activity against pathogenic bacterial and fungal strains.

KEYWORDS: *Cinnamomumzeylanicum*, stem bark, Volatile oil, Analysis, Microbiological activity

***Corresponding author:**

Prof. M. Ali

Phytochemistry Research Laboratory, Faculty of Pharmacy,
JamiaHamdard, Hamdard Nagar, New Delhi – 110 062, India

E Mail - maliphyto@gmail.com

INTRODUCTION

Cinnamomum zeylanicum Blume, syn. *C. verum* Presl. (Lauraceae), commonly known as dalchini or Ceylon cinnamon, is a moderate sized tree up to 16 m in height, native to Sri Lanka and southern India; distributed in several tropical countries, including China, Madagascar, West Indies, Malaysia and Java¹. Its bark is smooth, brown, thin, with a characteristic, strong pleasant odour and spicy burning taste. The bark is aromatic, astringent, expectorant and carminative; used to check nausea and vomiting and to treat stomach cramps, gastric irritation, dysentery, diarrhoea, neuralgia, rheumatism, toothache and paralysis of the tongue. It is one of the oldest and most valuable items in the spice trade used for flavouring liquids, in cakes, curries, tomato ketchup and sauces. The bark essential oil is utilized in food preparations, confectionery, perfumery, beverages, pharmaceuticals and dental preparations. It is perfect in topical applications, and with its pleasant scent, a perfect additive to creams, lotions, and soaps^{1,2}. It stimulates the uterine muscles and is used in difficult deliveries due to inadequate contractions. Cinnamon promotes regular and easy menstruation and is taken as a warming herb for cold conditions, often in combination with ginger (*Zingiber officinale*). It increases the circulation, especially to the fingers and toes and has been used to cure arthritis. It is also a traditional remedy for aching muscles and other symptoms of viral conditions such as colds and flu. Cinnamon tea is effective to stop bronchial asthma. Principally it is used as an aromatic to cover the unpleasant taste of other drugs. The cinnamon essential oil mainly contains cinnamaldehyde, eugenol, linalool and benzyl benzoate³⁻⁷. The oil possessed antifungal, antibacterial, antiparasitic, antispasmodic, antiflatulent, appetite stimulant, antioxidant, antidiarrhoeal, refrigerant and anthelmintic properties⁸⁻¹⁸. The present work describes analysis of the volatile and screening antimicrobial activity of the essential oil of the stem bark of *C. zeylanicum*.

MATERIALS AND METHODS

Plant material

The dried stem bark of *C. zeylanicum* was purchased from the Sangam Vihar Market, New Delhi. The plant material was identified by Dr. S.R. Mir, Department of Pharmacognosy and Phytochemistry, Jamia Humdard, New Delhi. A voucher specimen is preserved in the herbarium of the Department.

Isolation of the volatile oil

The dried stem bark powder (1 kg) of *C. zeylanicum* was hydrodistilled in an all glass Clevenger apparatus according to the method recommended in the British Pharmacopoeia, 1988. The colourless

volatile oil was dried over anhydrous sodium sulphate and stored at 4°C in the dark. The yield was 3.25 % based on the dried weight of the plant material.

GC analysis

Analytical GC was carried out by injection 01.µL of the leaf oil on a Varian 3300 gas chromatograph with FID detector fitted with silicone DB-I capillary column (30 m x 0.25 mm, film thickness 0.25 µm). GC operation condition was split mode, carrier gas was helium at a rate of 1.5 mL/min; temperature programme was from 80 to 225⁰ C (4⁰ C/min), injector temperature 280⁰ C and detector temperature 300⁰ C. Injection volume for all samples was 0.1 µl.

GC-MS analysis

GC-MS analysis was carried out by injection (0.1 µL) of the leaf oil on a QP-2000 instrument with a mass selective HP 597A detector fitted with Ulbon HR-1 capillary column (50 m x 0.25 mm, film thickness 0.25 µm). GC-MS operation condition was split mode, carrier gas helium at a flow rate of 1.5 mL/min; temperature programme 70-225⁰ C (10⁰ C/min), injector temperature 250⁰ C and detector temperature 280⁰ C. The mass spectrometry conditioned were as follows: ionization voltage, 70 eV; emission current, 40 mA; mass range 0 – 400 Da, ion source temperature, 200⁰ C.

Identification

The most constituents were identified by GC comparing their retention indices with those of authentic standard available in the laboratory or with the retention indices in close agreement with reference¹⁹. Further identification was achieved by analysis of GC/MS spectra. The fragmentation patterns of mass spectra were compared with those stored in the spectrometer data base using the NBS 54 KL and Wiley L-built libraries and with those reported in the literature^{19,20}.

Antimicrobial Activity

Test organisms and inoculums

Escherichia coli (NCTC-6571), *Staphylococcus aureus* (NCTC-10418) and *Bacillus subtilis* were obtained from the Division of Biotechnology, Faculty of Pharmacy, JamiaHamdard, New Delhi. The fungus strains *Aspergillusflavus*, *Aspergillusniger*, *Aspergillusfumigatus* and *Candida albicans* were procured from the Institute of Genomics and Integrative Biology (CSIR), New Delhi.

Antimicrobial standard

Tetracycline solution with specific activity of 50 µg/ml was prepared in DMSO solution (antibacterial). Fluconazole with specific activity of 50 µg/ml was prepared in DMSO solution (antifungal).

Media

Dehydrated nutrient agar media was prepared in distilled deionized water. The media (g/100 ml) was composed of peptone (5.1 g), sodium chloride (5.0 g), beef extract (1.5 g), yeast extract (1.5 g) and agar (1.5 g).

Preparation of media

Dehydrated nutrient agar medium (28 g) was accurately weighed and suspended in 1000 ml of distilled water in a conical flask. It was heated on a water bath to dissolve the medium completely. The conical flask containing the nutrient agar medium was plugged with the help of a non-absorbent cotton plug. Direct heating was avoided as it may lead to charring of the medium components and render it useless for the purpose.

Sterilization of Media

The conical flask containing the nutrient agar medium was plugged with the help of a non-absorbent cotton plug. The mouth of the conical flask and the cotton bung were properly covered with an aluminum foil. The medium was then sterilized by autoclaving at 15-lbs/in² pressure for 20 minutes.

Preparation of organisms

The test organisms were maintained on slants of medium and transferred to a fresh slant once a week. The slants were incubated at 37⁰C for 24 hours. Using 3 ml of saline solution, the organisms were washed from the agar slant on to a large agar surface (medium) and incubated for 24 hours at 37+2⁰C. The growth from the nutrient surface was washed using 50 ml of distilled water. A dilution factor was determined which gave 25 % light transmission at 530 nm. The amount of suspension to be added to each 100 ml agar or nutrient broth was determined by use of test plates or test broth. The test organisms were stored under refrigeration.

Temperature control

Thermostatic control is required in several stages of a microbial assay when culturing a micro-organism and preparing its inoculums and during inoculation in a plate assay.

Cup and plate method

A Previously liquefied and sterilized medium was poured into plastic petri-plates of 100 mm size. Required plates were prepared and kept for solidifying. Six holes were made in each plate with a stainless steel borer having 6 mm i.d. Different dilutions of the volatile oil of *C.zeylanicum* were made having concentration of 3 µl/ml, 5 µl/ml, 7 µl/ml and 9 µl/ml of solution. Tetracycline and fluconazole solutions were used as standards. The plates were labeled as Co (control), S (standard), X (*B. subtilis*), Y

(*S. aureus*), Z (*E. coli*) with four or five different holes, labeled as 3, 5, 7 and 9 for different concentrations. All dilutions were made in dimethyl sulphoxide (DMSO) solvent, which were used in experiment. Co was used for Control and S was used for Standard. Micropipette was used to deliver the solutions into the holes. The plates were then left for standing for 1 h for proper diffusion of the drug solutions. They were incubated for about 24 h at $32 \pm 2^\circ\text{C}$. After 24 h the plates were examined and the diameter of zones of inhibition was accurately measured. Antifungal activity was determined against *Aspergillusflavus*, *Aspergillusniger*, *Aspergillusfumigatus* and *Candida albicans* similar to antibacterial activity.

RESULTS AND DISCUSSION

The components of the volatile oil of the leaves of *C.zeylanicum*, their retention indices and percentage are listed in Table -1. The constituents are arranged in the order of their elution on Ulbon HR-1 capillary column. Analysis of the oil by GC-MS led to identification of twenty one components comprising 100% of the total volatile oil. The oil was characterized mainly by cinnamic aldehyde (91.5 %). There were thirteen monoterpenes (5.9 %) and ten of them were monoterpene hydrocarbons (5.1 %). The predominant monoterpene was *l*-limonene (3.1 %). There was three sesquiterpenes, viz. 3, 7-guaiadene, *cis*-calamenene and caryophyllene oxide, all of them were present in trace amounts. In addition to cinnamic aldehyde, the other aromatic components detected were benzaldehyde, *o*-cymene, eugenol, (E)-methyl cinnamate and benzyl benzoate, all of them occurred in lesser than 1 % yield. The oil was devoid of any aliphatic constituent.

The oil of the *C.zeylanicum* stem bark was examined for antibacterial activity against *E. coli*, *S. aureus* and *B. subtilis* and antifungal activity against *Aspergillusniger*, *A. fumigatus*, *A. flavus* and *Candida albicum*. The oil showed significant antimicrobial and antifungal activity in comparison to standards, Tetracycline and Fluconazole (Fig.1 and Fig. 2). The observations are recorded in the Tables 2 and 3.

Table 1: Percentage composition of volatile oil of the dried stem bark *C. zeylanicum*.

S.No.	Components	Retention Indices	Percentage
1	α -Pinene	925	0.2
2	Camphene	939	0.1
3	Sabinene	960	0.2
4	β -Pinene	964	0.2
5	Benzaldehyde	967	0.5
6	α -Phellandrene	991	0.1
7	<i>l</i> -Limonene	1017	3.1
8	<i>cis</i> -Ocimene	1019	0.1
9	<i>o</i> -Cymene	1023	0.8
10	Linalool	1077	0.3
11	Terpinenolene	1079	0.2
12	Γ -Terpenene	1083	0.1
13	Thujyl alcohol	1125	0.2
14	Terpinen-4-ol	1129	0.3
15	Cinnamicaldehyde	1210	91.5
16	Eugenol	1326	0.5
17	3,7-Guaiadene	1430	0.3
18	<i>cis</i> -Calamenene	1451	0.4
19	(E) –Methyl cinnamate	1507	0.5
20	Caryophyllene oxide	1520	0.2
21	Benzyl benzoate	1866	0.2

Table 2: Antibacterial activity of the stem bark of *C. zeylanicum*..

Sample conc. (μ l/ml)	Zone of inhibition (mm) <i>S. aureus</i>	Zone of inhibition (mm) <i>E. coli</i>	Zone of inhibition (mm) <i>B. subtilis</i>
3 (A)	12	10	12
5 (B)	17	14	14
7 (C)	18	15	16
9 (D)	18	17	27
50 (Co)	---	---	---
50 (S)	11	16	27

Table 3: Antifungal activity of the stem bark of *C. zeylanicum*.

Sample conc. ($\mu\text{l/ml}$)	Zone of inhibition (mm) <i>A. niger</i>	Zone of inhibition (mm) <i>A. fumigatus</i>	Zone of inhibition (mm) <i>A. flavus</i>	Zone of inhibition (mm) <i>C. albicum</i>
3 (A)	22	25	27	23
5 (B)	27	27	28	25
7 (C)	28	33	35	26
9 (D)	33	38	37	36
50 (Co)	---	----	---	---
50(S)	22	12	14	30




Fig. 1: Antibacterial activity of the stem bark of *C. zeylanicum*




Fig. 2: Antifungal activity of the stem bark of *C. zeylanicum*

CONCLUSION

The volatile oil of *C. zeylanicum* was consisted mainly of mainly by cinnamic aldehyde (91.5 %). There were thirteen monoterpenes (5.9 %) and three sesquiterpenes. It showed significant antimicrobial activity.

ACKNOWLEDGEMENT

The authors are thankful to the Head, SAIF, CDRI, Lucknow, for recording GC and CG-MS analysis of the volatile oil.

REFERENCES

1. Anonymous; .The Wealth of India., New Delhi, RawMaterials,Publication and Information Directorate, CSIR, New Delhi. 1992;3: 582 – 589.
2. KirtikarKR ,Basu BD. Indian Medicinal Plants. India, Sri Satguru Publication, Delhi. 2000; 2966-2968.
3. Chericoni S, PrietoJM ,Iacopini P, Cioni P, Morelli I. In Vitro Activity of the Essential Oil of *Cinnamomumzeylanicum* and Eugenol in Peroxynitrite-Induced Oxidative Processes, J.Agric. Food Chem., 2005; 53 (12): 4762–4765.

4. Arora P , Nanda A, Karan M. GC-MS Profile of Volatile oils of *Cinnamomum Zeylanicum* Blume and *Ocimum kilimandscharicum* Baker ex Gurke . Int. J. Pharm. Sci. Rev. Res. 2013; 19(2): 124-126.
5. Nkanwen ERS, Awouafack MD, Jules J, Bankeu K, Wabo HK , Mustafa SAA, Ali MS, Lamshöft M, Choudhary MI , Michael Spittler M, Tane P. Constituents from the stem bark of *Cinnamomum zeylanicum* Welw. (Lauraceae) and their inhibitory activity toward *Plasmodium falciparum* Enoyl-ACP reductase enzyme. Rec. Nat. Prod. 2013; 7(4): 296-301.
6. Senanayake TU, Lee MH, Wills RBH. Volatile constituents of Cinnamon (*Cinnamomum zeylanicum*). Oils, J. Agric. Food. Chem. 1978; 26: 822-824.
7. Al-Bayati FA, Mohammed MJ. Isolation, identification, and purification of cinnamaldehyde from *Cinnamomum zeylanicum* bark oil. An antibacterial study. Pharmaceutical Biology. 2009, 47 (1): 61-66.
8. Mastura M, Azah MAN, Khozirah S, Mawardi R, Manaf AA, Anticandidal and antidermatophytic activity of *Cinnamomum* species essential oils. Cytobios. 1999; 98: 17-23.
9. Štefan Faix, Zita Faixová, Iveta Plachá, Juraj Koppel, Effect of *Cinnamomum zeylanicum* Essential Oil on Antioxidative Status in Broiler Chickens. Acta Vet. Brno 2009; 78: 411-417.
10. Fabian D, Sabol M, Domaracka K, Bujnakova D. Essential oils-their antimicrobial activity against *Escherichia coli* and effect on intestinal cell viability. Toxicol In Vitro. 2006; 20: 1435-1445
11. Tomaino A, Cimino F, Zimbalatti V, Venuti V, Sulfaro V, De Pusquale A, Saiha A. Influence of heating on antioxidant activity and the chemical composition of some spice essential oils. Food Chem. 2005; 89: 549-554.
12. S. Elumalai, R. Kesavan, S. Ramganesan, R. Murugasen. Isolation, purification and identification of the antidiabetic components from *Cinnamomum zeylanicum* and *Cinnamomum cassia* bark oil extracts. Curr. Bot. 2011; 2(2): 12-17.
13. Rathi B, Bodhankar S, Mohan V, Thakurdesai P. Ameliorative effects of a polyphenolic fraction of *Cinnamomum zeylanicum* L. bark in animal models of inflammation and arthritis. Sci Pharm. 2013; 81(2): 567-589.
14. Nyadjeu P, Nguielefack-Mbuyo EP, Atsamo AD, Nguielefack TB, Dongmo AB, Kamanyi A. Acute and chronic antihypertensive effects of *Cinnamomum zeylanicum* stem bark methanol

- extract in L-NAME-induced hypertensive rats. BMC Complement Altern Med. 2013; Jan 31;13:27. doi: 10.1186/1472-6882-13-27.
15. Nyadjeu P, Dongmo A, Nguelefack TB, Kamanyi A. Antihypertensive and vasorelaxant effects of *Cinnamomumzeylanicum* stem bark aqueous extract in rats. J Complement Integr Med. 2011; Jan;8. doi: 10.2202/1553-3840.1490.
16. Unlu M, Ergene E, Unlu GV, Zeytinoglu HS, Vural N. Composition, antimicrobial activity and in vitro cytotoxicity of essential oil from *Cinnamomumzeylanicum* Blume (Lauraceae). Food Chem Toxicol. 2010; 48(11):3274-80, doi: 10.1016/j.fct.2010.09.001.
17. Chase CK, McQueen CE. Cinnamon in diabetes mellitus. Am J Health Syst Pharm. 2007; 64(10):1033-1035.
18. Kanuri G, Weber S, Volynets V, Spruss A, Bischoff SC, Bergheim I. Cinnamon extract protects against acute alcohol-induced liver steatosis in mice. J Nutr. 2009; 139(3):482-487. doi: 10.3945/jn.108.100495.
19. Adam RP. Identification of essential oil components by gas chromatography/quadrupole mass spectrometry. Carol Stream IL, USA, Allured. 2001.
20. Ali M. Techniques in terpenoid identification. Birla Publishers, Delhi. 2001;4-51.