

International Journal of Research in Pharmacy and Science

Arsenic in Drinking Water: Affect the Gene Expression Related to type II Diabetes

Sagar M. S¹, Rashmi N. G* and Gagana Aradhya¹

¹Department of Pharmaceutical Analysis, Sri Adichunchanagiri College of Pharmacy,
Adichunchanagiri University, B.G. Nagara, Karnataka, India.
E Mail - sagarmspharma96@gmail.com, Gaganaaradhya05@gmail.com

ABSTRACT

Chronic exposure to high concentrations of arsenic in drinking water is associated with an increased risk for developing type II diabetes. The present revision focusses on the association of arsenic in water more than above normal, which might be affecting on the expression of genes related to type II diabetes with *in vivo* experimental evidence. We identified a few *in vivo* studies reported that arsenic interfered with transcription factors involved in insulin-related gene expression in pancreatic beta-cells arsenic decrease insulin synthesis, secretion and reduce the expression of antioxidant enzymes and in the liver whereas arsenic disturbs glucose productions by alteration of gluconeogenesis and abnormal proliferation. Currently available *in vivo* experimental study evidence shows that, the association between high and chronic exposure to arsenic in drinking water developing type II diabetes. But the effect of exposure to low or moderate levels of arsenic on risk in diabetes is unclear. An increase in the prevalence of diabetes mellitus has been observed among residents of highly arsenic contaminated areas.

KEYWORDS: Arsenic, Drinking water, Type II diabetes, Gene expression, Pancreatic beta-cells, Gluconeogenesis.

*Corresponding Author:

Rashmi N. G*

Department of Pharmaceutical Analysis,
Sri Adichunchanagiri College of Pharmacy, Adichunchanagiri University,
B.G. Nagara, Karnataka, India.

E Mail - ng.rashmi85@gmail.com

1. INTRODUCTION

Type 2 diabetes mellitus is a metabolic disorder characterized by hyperglycaemias, insulin resistance in peripheral tissues, and altered insulin secretory capacity of pancreatic β -cells. Type 2 diabetes accounts for 90-95% of all cases of diabetes and is a major public health problem worldwide¹. Established risks factors of type 2 diabetes include older age, obesity, physical inactivity, family history, and genetic polymorphisms.

In addition, environmental toxicants, including arsenic, have been suggested to play an etiologic role in diabetes development². Arsenic is a recognized toxicant and carcinogen. Non-occupational exposure occurs mainly through drinking water, affecting millions of people worldwide. Exposure to levels of arsenic in drinking water well above 100 ppb has been associated with an increased risk of type 2 diabetes. Our objective was to perform a systematic review of the experimental studies (in vitro or in vivo) to synthesize available information on plausible mechanisms for gene expression which is related to type 2 diabetes.

The potential mechanisms proposed for arsenic-induced type 2 diabetes include, phosphorus substitution, high affinity for sulfhydryl groups, increased oxidative stress and interference with gene expression³. Arsenic has been shown to alter the expression of genes related to the maintenance of glucose homeostasis in tissues associated with the pathogenesis of the disease. Thus, arsenic could lead to diabetes by impairing the expression of a wide range of genes involved in pathways that maintain this homeostasis.

Pancreatic islets, adipose tissue, muscle and the liver, regulate glucose blood levels. Pancreatic β -cells secrete insulin in response to an increase in blood glucose levels. This hormone promotes the entrance of glucose into adipocytes and muscle cells via the glucose transporter GLUT4, and inhibits hepatic glucose production. The liver is responsible for gluconeogenesis, which maintains acceptable blood glucose levels when dietary glucose intake is insufficient. In addition, insulin stimulates cell growth and differentiation and promotes the storage of substrates in fat, liver and muscle by stimulating lipogenesis, glycogen and protein synthesis, and by inhibiting lipolysis, glycogenolysis and protein breakdown⁴. In the present review, we discuss the expression of genes involved in type 2 diabetes that are known to be affected by arsenic, identifying possible mechanisms or pathways by which arsenic could lead to or promote diabetes.

2. DEOXYRIBONUCLEIC ACID (DNA) METHYLATION

DNA methylation, a major regulator of epigenetic modifications has been shown to alter the expression of genes that are involved in aspects of glucose metabolism such as glucose intolerance, insulin resistance, β -cell dysfunction and other conditions, and it ultimately leads to the pathogenesis of type 2 diabetes mellitus (T2DM). Current evidences indicate an association of DNA methylation with T2DM. This review provides an overview of how various factors play crucial roles in T2DM pathogenesis and how DNA methylation interacts with these factors. Additionally, an update on current techniques of DNA methylation analysis with their pros and cons is provided as a basis for the adoption of suitable techniques in future DNA methylation research towards better management of T2DM. To elucidate the mechanistic relationship between vital environmental factors and the development of T2DM, a better understanding of the changes in gene expression associated with DNA methylation at the molecular level is still needed.⁵⁻¹⁰

Arsenic influences the sensitivity of insulin by changing the expression pattern of the genes involved in the insulin resistance and shifts away the cells from differentiation to proliferation status¹¹. Many studies have shown the role of oxidative stress and inflammatory response as the potent key mediator of arsenic induced toxicity¹²⁻¹⁷. Exposure to a low level of arsenite changes the gene expression of inflammatory cytokines such as interleukin 6 as well as genes involved in oxidative stress response leading to a decrease in insulin secretion, pancreatic islets dysfunction and changes in gluconeogenesis¹⁸⁻²³. The most extensively studied epigenetic mechanisms are the alteration of DNA methylation pattern and histone post-translation modifications, which in turn regulate the expression of genes at the level of transcriptional. Epigenetic changes such as DNA methylation alterations have been considered as a potential underlying mechanism in the pathophysiology of many diseases including Diabetes Mellitus²⁴⁻²⁷. In DNA methylation critical genes which are involved in pancreatic development, insulin secretion and glucose sensing including insulin 2 (Ins2), pancreatic and duodenal homeobox 1 (Pdx1), peroxisome proliferator-activated receptor gamma (PPAR γ) and glucose transporter 2 (Glut2), in order to evaluate DNA methylation alterations as a potent underlying mechanism of arsenic induced pancreatic dysfunction and diabetes development.²⁸⁻³⁰

2.1 Deoxyribonucleic Acid (DNA) Methylation Analysis

Fazlullah khan²⁴ evaluated for the methylation status of the Pdx1, PPAR γ , Ins2 and Glut2 genes after 6 days of exposure. The genomic DNA was isolated from 250 islets cells and was assessed for the amount, purity and integrity of DNA using ultraviolet-visible (UV) spectrophotometer and gel

electrophoresis. The bisulfite conversion procedures were performed as per the EZ DNA Methylation kit (Zymo Research Corporation, USA). Two separate bisulfite conversion treatment were performed for each DNA sample.

3. Messenger Ribonucleic Acid (mRNA)

Arsenic-induced alteration in mRNA insulin expression and secretion (pancreatic β -cell dysfunction) fits the model where insulin resistance alone is not enough to develop type 2 diabetes. Instead, both insulin resistance and insulin deficiency are needed in conjunction.

The mRNAs that were found significantly reduced were: (a) the transcription factor required for adipogenesis, peroxisome proliferative-activated receptor gamma (PPAR γ); (b) the adipocyte selective fatty acid-binding protein (AP2) (induced by PPAR γ); (c) the transcription factor CCAAT-enhancer binding protein alpha (C/EBP α) that works in conjunction with PPAR γ to induce the expression of most fat cell-specific genes; and (d) genes involved in cell cycle regulation, like p21Cip1/Waf1 and p27Kip1^(31,32). Arsenic exposure causes significant downregulation of mRNA expression of immune response related genes in most of the experimental groups^{33,34} (Table 1)

Table 1: Messenger Ribonucleic acid (mRNA) expression of immune response related to genes

Sl No		Time	Temperature	Dose	Effect		
01	Dose dependent response [single exposure]	24h	-80°C	LD24	Downregulated NFkb signalling genes		
				MD24	Downregulated expression of inhibitor Nfkbia		
				HD24			
02	Time dependent response [24h vs 48h]	24h vs 48h	-80°C	LD24	6↓↑	5↑↑	One is unique to LD24
				LD48			
03	Effects of dose frequency	24h	-80°C	RD	Expression of Chuk was downregulated		
				RDCE			

LD24 - Low dose

MD24 - Medium dose

HD24 - High dose

RD - Repeated dose

RDCE-Repeated dose cessation

↓↑- Upregulated or Downregulated

↑↑- Upregulated

4. CONCLUSION

Millions of people all over the world are constantly exposed to iAs through the consumption of contaminated drinking water. Different mechanisms have been suggested for arsenic-induced diabetes. Our results showed that arsenic exposure alters DNA methylation status of key insulin-related genes. Also, we found that sodium arsenite impacts the expression of other key regulatory genes involved in insulin metabolism and therefore could result in the impairment of islets function. There is a possibility of the involvement of other epigenetic regulatory mechanisms such as histone post-translational modifications or microRNA dysregulation that remained to be explored in the future. Additionally, a further longitudinal in vivo study is suggested to establish the sodium arsenite-induced methylation changes as the underlying mechanism involved in islet dysfunction. Our results reinforce the importance of future research on arsenic-induced epigenetic modifications aiming at finding therapeutic or prognostic factors.

This study demonstrated that exposure to arsenic may have substantial effects on gastrointestinal health, possibly through the gene expression of the various immune response-related pathways. While the potential health effects of long-term arsenic exposure are well known, it is important to address the lack of knowledge of the impact of low doses of arsenic on gastrointestinal health. These data show that low-level arsenic exposure may have a dramatic impact on immune system preparedness, even after a single exposure.

The effects of arsenic both altering pancreatic β -cell function and glucose uptake in peripheral tissues, which resemble type 2 diabetes, suggest that these mechanisms play an important role in the increased incidence of type 2 diabetes in chronically exposed to chronic inorganic arsenic(iAs)

5. REFERENCE:

1. DeFronzo RA, Ferrannini E, Groop L, et al. Type 2 diabetes mellitus. *Nature reviews Disease primers*. 2015 Jul 23;1(1):1-22.
2. DeFronzo RA. Pathogenesis of type 2 diabetes mellitus. *Medical Clinics*. 2004 Jul 1;88(4):787-835.
3. Akash MS, Rehman K, Chen S. Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus. *Journal of cellular biochemistry*. 2013 Mar;114(3):525-31.
4. Saltiel AR, Kahn CR. Insulin signalling and the regulation of glucose and lipid metabolism. *Nature*. 2001 Dec;414(6865):799-806.

5. Alam F, Islam A, Hua Gan S, Mohamed M, Haryo Sasongko T. DNA methylation: an epigenetic insight into type 2 diabetes mellitus. *Current pharmaceutical design*. 2016 Aug 1;22(28):4398-419.
6. Tseng CH, Tai TY, Chong CK, et.al, Long-term arsenic exposure and incidence of non-insulin-dependent diabetes mellitus: a cohort study in arseniasis-hyperendemic villages in Taiwan. *Environmental health perspectives*. 2000 Sep;108(9):847-51.
7. Smeester L, Rager JE, Bailey KA, et.al, Epigenetic changes in individuals with arsenicosis. *Chemical research in toxicology*. 2011 Feb 18;24(2):165-7.
8. Abdul KS, Jayasinghe SS, Chandana EP, Jayasumana C, De Silva PM. Arsenic and human health effects: A review. *Environmental toxicology and pharmacology*. 2015 Nov 1;40(3):828-46.
9. Paul DS, Harmon AW, Devesa V, Thomas DJ, Stýblo M. Molecular mechanisms of the diabetogenic effects of arsenic: inhibition of insulin signaling by arsenite and methylarsonous acid. *Environmental health perspectives*. 2007 May;115(5):734-42.
10. Fu J, Woods CG, Yehuda-Shnaidman E, et. al, Low-level arsenic impairs glucose-stimulated insulin secretion in pancreatic beta cells: involvement of cellular adaptive response to oxidative stress. *Environmental health perspectives*. 2010 Jun;118(6):864-70.
11. Cavaghan MK, Ehrmann DA, Polonsky KS. Interactions between insulin resistance and insulin secretion in the development of glucose intolerance. *The Journal of clinical investigation*. 2000 Aug 1;106(3):329-33.
12. Shirayama M, Seth M, Lee HC, et. al, piRNAs initiate an epigenetic memory of nonself RNA in the *C. elegans* germline. *Cell*. 2012 Jul 6;150(1):65-77.
13. Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, Reese SE, Markunas CA, Richmond RC, Xu CJ, Küpers LK. DNA methylation in newborns and maternal smoking in pregnancy: genome-wide consortium meta-analysis. *The American Journal of Human Genetics*. 2016 Apr 7;98(4):680-96.
14. Broberg K, Ahmed S, Engström K, et. al, Arsenic exposure in early pregnancy alters genome-wide DNA methylation in cord blood, particularly in boys. *Journal of developmental origins of health and disease*. 2014 Aug;5(4):288-98.
15. Michailidi C, Hayashi M, Datta S, et. al, Involvement of epigenetics and EMT-related miRNA in arsenic-induced neoplastic transformation and their potential clinical use. *Cancer prevention research*. 2015 Mar 1;8(3):208-21.
16. Kuo CC, Howard BV, Umans JG, et. al, Arsenic exposure, arsenic metabolism, and incident diabetes in the strong heart study. *Diabetes care*. 2015 Apr 1;38(4):620-7.

17. Liu S, Guo X, Wu B, et. al, Arsenic induces diabetic effects through beta-cell dysfunction and increased gluconeogenesis in mice. *Scientific reports*. 2014 Nov 4;4:6894.
18. Douillet C, Currier J, Saunders J, et. al, Methylated trivalent arsenicals are potent inhibitors of glucose stimulated insulin secretion by murine pancreatic islets. *Toxicology and applied pharmacology*. 2013 Feb 15;267(1):11-5.
19. Kaushal A, Zhang H, Karmaus WJ, et. al, Genome-wide DNA methylation at birth in relation to in utero arsenic exposure and the associated health in later life. *Environmental Health*. 2017 Dec;16(1):1-1.
20. Majumdar S, Chanda S, Ganguli B, et. al, Arsenic exposure induces genomic hypermethylation. *Environmental toxicology*. 2010 Jun;25(3):315-8.
21. Kile ML, Houseman EA, Baccarelli AA, et. al, Effect of prenatal arsenic exposure on DNA methylation and leukocyte subpopulations in cord blood. *Epigenetics*. 2014 May 6;9(5):774-82.
22. Houseman EA, Accomando WP, Koestler DC, et. al, DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC bioinformatics*. 2012 Dec 1;13(1):86.
23. Reichard JF, Schneckenger M, Puga A. Long term low-dose arsenic exposure induces loss of DNA methylation. *Biochemical and biophysical research communications*. 2007 Jan 5;352(1):188-92.
24. Khan F, Hodjat M, Rahimifard M, et. al, Assessment of arsenic-induced modifications in the DNA methylation of insulin-related genes in rat pancreatic islets. *Ecotoxicology and Environmental Safety*. 2020 Sep 15;201:110802.
25. Weber M, Hellmann I, Stadler MB, et. al, Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome. *Nature genetics*. 2007 Apr;39(4):457-66.
26. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nature Reviews Genetics*. 2012 Jul;13(7):484-92.
27. Moore LD, Le T, Fan G. DNA methylation and its basic function. *Neuropsychopharmacology*. 2013 Jan;38(1):23-38.
28. Wauson EM, Langan AS, Vorce RL. Sodium arsenite inhibits and reverses expression of adipogenic and fat cell-specific genes during in vitro adipogenesis. *Toxicological Sciences*. 2002 Feb 1;65(2):211-9.
29. Chai CY, Huang YC, Hung WC, Kang WY, Chen WT. Arsenic salts induced autophagic cell death and hypermethylation of DAPK promoter in SV-40 immortalized human uroepithelial cells. *Toxicology letters*. 2007 Aug 30;173(1):48-56.

30. Yoshida T, Yamauchi H, Sun GF. Chronic health effects in people exposed to arsenic via the drinking water: dose–response relationships in review. *Toxicology and applied pharmacology*. 2004 Aug 1;198(3):243-52.
31. Chen YW, Yang CY, et. al, Heavy metals, islet function and diabetes development. *Islets*. 2009 Nov 1;1(3):169-76.
32. Andrew AS, Burgess JL, Meza MM, et. al, Arsenic exposure is associated with decreased DNA repair in vitro and in individuals exposed to drinking water arsenic. *Environmental health perspectives*. 2006 Aug;114(8):1193-8.
33. Andrew AS, Burgess JL, Meza MM, et. al, Arsenic exposure is associated with decreased DNA repair in vitro and in individuals exposed to drinking water arsenic. *Environmental health perspectives*. 2006 Aug;114(8):1193-8.
34. Arnold MG, Gokulan K, Doerge DR, Vanlandingham M, Cerniglia CE, Khare S. A single or short time repeated arsenic oral exposure in mice impacts mRNA expression for signaling and immunity related genes in the gut. *Food and Chemical Toxicology*. 2019 Oct 1;132:110597.