

Review article

Available online www.ijrpsonline.com

International Journal of Research in Pharmacy and Science

Review on Leonotis nepetifolia Linn

Jeyaraman AmuthaIswarya Devi, Murugan Vannithurai*and Velayutham Mani Mala

Department of Pharmaceutical Chemistry, ArulmiguKalasalingam College of Pharmacy, Anand Nagar, Krishnankoil-626126, Srivilliputtur (via) Tamil Nadu, India. E-mail id: <u>vanishdathav@gmail.com</u>

ABSTRACT

The plant of *Leonotis nepetifolia* Linn in India, belongs to family *Lamiaceae. Leonotis nepetifolia* is commonly called 'kilp dagga' or 'lion ear', 'shandilay' and 'bradi-bita'. It is native to tropical Africa and southern India. It grows to a height of 3 metres and striking lipped flowers that are most commonly orange but can vary to red, white, and purple. It has drooping dark green, very soft serrated leaves that can grow up to 10 centimetres wide. The plant stems emanates from a thick wood base. Sunbirds and ants are attracted to the flowers. It has been found growing on road sides, rubbish heaps or waste land. The cultivation of medicinal plants is the key to meet the raw material needs of the industry based on the medicinal plant products. Phytochemical investigation shows the presence of alkaloids, protein & amino acids, flavonoids, terpenoids, phenolics, glycosides, carbohydrate, saponins, steroids, tannins and fixed oil. Gas Chromatography – Mass Spectoscopy analysis of showed several active components. A review lowed formed on various parts of this plant are anti-inflammatory activity, anti-bacterial, anti-diarrheal, Healing, antiviral, antifungal, anti-giardial, analgesic, anti-diabetic, anti-thelmintic, antioxidant, antimicrobial and cytotoxic activity.

KEYWORDS: Lamiaceae, Leonotisnepetifolia, Lion's ear, Ranabheri.

*Corresponding author

Murugan Vannithurai

Department of Pharmaceutical Chemistry

ArulmiguKalasalingam College of Pharmacy,

Srivilliputtur (via) Tamil Nadu, India.

E-mail id: vanishdathav@gmail.com

INTRODUCTION

Herbal medicine (also Herbalism) is the study of the botany and use of medicinal plants. Plants have been the basis for medical treatments through much of human history, and such traditional medicine is still widely practiced today. The scope of herbal medicine is sometimes extended to include fungal and bee products, as well as minerals, shells and certain animal parts. Herbal medicine is also called phytomedicine or phytotherapy. Para herbalism is the pseudoscientific use of extracts of plant or animal origin as supposed medicines or health-promoting agents.¹⁸

Medicinal plants are the nature's gift to human beings to help them pursue a disease-free healthy life. The world's cultures have an extensive knowledge of herbal medicine. Plants are a valuable source of a wide range of secondary metabolites, which are used as pharmaceuticals, agrochemicals, flavours, fragrances, colours, bio pesticides and food additives.¹⁸

The genus *Leonotis* has 12 species widely distributed in pan tropics and is represented by one species, *Leonotis nepetifolia* in India. It belongs to family *Lamiaceae*. *Leonotis nepetifolia* is important medicinal plant of repute in Indian traditional systems of medicine such as Ayurveda, Unani and Siddha. Medicinal uses of the plant are reported in Madagascar, Brazil, Canada, Kenya and many African Countries to treat diseases, rheumatism, dysmenorrhoea, bronchial asthma, fever and diarrhoea^{9,3}.

PLANT DESCRIPTION

Leonotis nepetifolia Linn plant belong to genus *Leonotis* and family *Lamiaceae*. Commonly it's know "Lion ear" and hindis "Baraguma". It is a medium sized deciduous plant and very vigorous in growth. *Leonotis nepetifolia* is otherwise called as 'Ranabheri' in tamil. It is one of the most traditional system of medicine in Ayurvedic and Siddha.^{9,3}

Botanical name	:Leonotis nepetifoliaLinn.
Family	:Lamiaceae
Kingdom	:Plantae
Order	:Lamiales
Subfamily	:Lamioidae
Genus	:Leonotis
Species	:Leonotisnepetifolia

Domain	:Eukaryota
Phylum	:Spermatophyta
Subphylum	:Angiospermae
Class	:Dicotyledonae
Common name	:Annual lion's ear, lion's ear, flor de mundo and chrishmas candlestick
Synonyms	:L.globosusMoench, L.nepetifoliusMill andPhlomisnepetifoliaLinn.



Fig. 1: Whole plant of LeonotisnepetifoliaLinn.

VERNACULAR NAMES

Tamil: Ranabheri.

Sanskrit: Granthi and Granthika.

Kannada: Deepashoole.

Telugu: BaeriandBeri.

Gujarati: Lalguma and Mahadrona.

English: Bald bush and Bald head.

Hindi: Baraguma and Hejurchel.

MEDICINAL USES³

- 1. *Leonotis* may be involved in the anti-nociceptive, anti-inflammatory and anti-diabetic effect of the plant's extract.
- 2. Which has been used to treat bronchial asthma, diarrhoea, fever, influenza, malaria and analgesic.
- 3. This herb in control of painful, arthritic, and other inflammatory conditions.
- 4. It is used to type2 diabetes mellitus.

- 5. It is also anti-asthmatic and has anti-diarrhea properties.
- 6. They are used for heart conditions associated with anxiety and tension.

FOLK MEDICINAL USES³

- 1. The fresh stem juice is an infusion drunk for 'blood impurity'. The infusions made from flowers and seeds, leaves or stems are widely used as tonics for tuberculosis, jaundice, muscular cramps, high blood pressure, diabetes, viral hepatitis, dysentery, and diarrhoea.
- 2. The leaf tea has a hypnotic effect, is diuretic and relieves headache.
- 3. The leaf and stem decoction or inhalations have been used internally for cough, common cold, bronchitis, wound healing and asthma.
- 4. Tea made from the whole plant is used for arthritis, piles, bladder and kidney disorder, obesity, cancer and rheumatism.
- 5. The leaves and stems decoction are applied topically as a treatment for eczema, skin infections and itchiness.
- 6. The leaves, roots and bark are widely used as an emetic for snakebites, bee and scorpion stings.

OTHER USES³

1. The seeds are rich in a fatty oil resembling olive oil

PHYSICOCHEMICAL INVESTIGATION

Physicochemical screening of *Leonotis nepetifolia* revealed that, it possess good physicochemical parameters such as total ash, acid insoluble ash, water soluble ash, loss on drying, swelling index and foaming index. The extractive values such as alcohol soluble extract, water soluble extract, pet ether, benzene, chloroform, ethyl acetate, methanol and aqueous extract are also determined.

S. No	Parameters	Observation of leaves	Observation of root
1.	Total ash	5.88 % w/w	4.083 % w/w
2.	Acid insoluble ash	1.003 % w/w	0.7067% w/w
3.	Water soluble ash	4.1 % w/w	3.15% w/w
4.	Foreign organic matter	0.88 % w/w	1.85% w/w
5.	Moisture content	7.8% w/w	5.1% w/w
6.	Foaming index	Less than 100	Less than 100

 Table 1: Physicochemical parameters of Leonotisnepetifolia¹⁶

S. No	Parameters	Observation of leaves	Observation of root
1.	Alcohol soluble extractive	29.2 % w/w	21.83 % w/w
2.	Water soluble extractive	27.97 % w/w	28.39 % w/w
3.	Pet ether	4.48 % w/w	0.05 % w/w
4.	Benzene	4.17 % w/w	0.48 % w/w
5.	Chloroform	1.63 % w/w	2.43 % w/w
6.	Ethyl acetate	1.47 % w/w	0.16 % w/w
7.	Methanol	21.4 % w/w	22.53 % w/w
8.	Aqueous	15.04 % w/w	18.25 % w/w

 Table 2: Extractive values of Leonotisnepetifolia
 6,16

PRELIMINARY PHYTOCHEMICAL INVESTIGATION

The petroleum ether, ethyl acetate, chloroform, methanol and aqueous extracts were subjected to preliminary phytochemical screening for detection of various plant constituents present. They are individually performed using different qualitative tests for alkaloids, carbohydrates, flavonoids, glycosides, protein & amino acids, saponins, fixed oils, terpenoids, phenolics, tannins and steroids.

S. No	Type of Phyto Constituents	Petroleum ether extract	Ethyl acetate Extract	Chloroform extract	Methanol extract	Aqueous extract
1.	Alkaloids	-	-	+	+	-
2.	Carbohydrates	-	-	_	+	+
3.	Flavonoids	-	+	-	+	+
4.	Glycosides	-	-	+	+	+
5.	Protein & amino acids	-	-	-	-	+
6.	Saponins	-	-	-	-	+
7.	Fixed oil	+	-	-	-	-
8.	Terpenoids	+	-	+	-	-
9.	Phenolics	-	-	-	+	+
10.	Steroids	+	+	+	-	-
11.	Tannins	+	-	-	-	-

 Table 3: Phytochemical Investigation of leaves of Leonotisnepetifolia

S. No	Type of phyto Constituents	Petroleum ether extract	Ethyl acetate extract	Chloroform extract	Methanol extract	Aqueous extract
1.	Alkaloids	-	-	-	-	-
2.	Carbohydrates	-	+	-	+	+
3.	Flavonoids	-	+	-	+	+
4.	Glycosides	-	-	+	+	+
5.	Protein & amino acids	-	-	-	+	+
6.	Saponins	-	-	-	+	+
7.	Fixed oil	+	-	+	-	-
8.	Terpenoids	+	-	+	-	-
9.	Phenolics	-	+	-	+	+
10.	Steroids	+	+	+	-	-
11.	Tannins	-	+	-	+	+

 Table 4: Phytochemical Investigation of root of Leonotisnepetifolia

PHARMACOLOGICAL ACTIVITY

Table: 5Ethnopharmacology of Leonotis nepetifolia Linn.

S.	Plant part	Solvent used for	Activity	Author name
No		extraction		
1	Leaves	Ethanolic extract	Anti-bacterial, anti- diarrheal,	Darley Maria Oliveira et
			healing and antiviral	al., 2015
2	Leaves	n- Hexane and	Antibacterial, antifungal and	UdayaPrakashet al., 2013
		Methanolic extract	antiviral	
3	Flower	Methanolic extract	Antifungal	Wagara IN et al., 2015
4	Root	Methanolic extract	Anti-giardial	Ndukui James Gakunga
				et al., 2013
5	Whole plant	Ethanolic extract	Analgesic and anti-diabetic	Neeraj Sethiyaet al., 2011
6	Whole plant	Ethanol and aqueous	Anti-thelmintic	Reshmi Pushpanet al.,
		extract		2017
7	Aerial part	Methanol and Acetone	Antibacterial and cytotoxic	Sarath Chandirianet al.,
		extract		2017
8	Leaves, flower	Ethanolic extract	Antimicrobial and cytotoxic	Tiziana Fornariet al.,
	and stem		activity	2012
9	Leaves, flower	n-Hexane, methylene	Anti-inflammatory	Ana Paula de Oliveira et
	and stem	dichloride and		al., 2015
		methanolic extract		
10	No specified	Methanolic extract	Antioxidant	Usharani Veerabadranet
				al., 2013

GAS CHROMATOGRAPHY - MASS SPECTROSCOPY ANALYSIS

GC-MS analysis of the fixed oils from the leaves of specimens of *L. nepetifolia*showed 16 bioactive compounds were identified such as Methyl linoleate (46.98%) was the majority compound. Among these, two pairs of isomers, propanoic acid 2-methyl-3-hydroxy-2,4,4-trimethylpentyl ester (31.97%) and propanoic acid 2-methyl-2,2-dimethyl-1-(2-hydroxy-1-methylethyl)-propyl ester (22.78%). This compound like methyl laurate (0.38%), methyl myristate (0.64%), phytol (0.49%), cycloartenol (1.93%), methyl palmitate(13.67%), β -amyrim (2.58%), y-undecanolide (1.32%), y-decanolide(7.67%), 9,12-octadecadienoic acid methyl ester(9.21%), 6-octadecynoic acidmethyl ester (6.49%),methyl stearate(2.79%), arachidic acid methyl ester(0.74%), docosanoicacifd methyl ester(0.45%), squqlene(0.62%), stigmasterol (1.62%), stigmast-5-en-3 β -ol(1.11%), stigmast-7-en-3 β -ol(0.95%), pentacosane(0.48%).⁷

GC–MS study was analysed essential oil from*Leonotis nepetifolia*oil was obtained in 0.004% w/w yield. A total of 36 compounds constituting 94.80% of the oil were identified. The oil was characterized by a high percentage of sesquiterpenes (75.32%). The major compounds were germacrene D (37.16%),germacrene–D-4-ol (6.02 %,) and *cis*-ocimene (4.79%).

RESULT AND DISCUSSION

The survey of literature revealed that the *Leonotis nepetifolia* having effective pharmacological activities such as anti-inflammatory activity, anti-bacterial, anti-diarrheal, healing, anti-viral, anti-fungal, anti-giardial, analgesic, anti-diabetic, anti-thelmintic, anti-oxidant, anti-microbial and cytotoxic activity. It having important medicinal phytochemical compounds such asalkaloids (leonuride), iridoid glycoside (leonurin and leonuridine), diterpenoides (leocardin), flavonoids (ratin, quercetin, hyperoside, apigenin), volatile oil, tannins, vitamin A. The results from this review are quite promising for the use of *Leonotis nepetifolia* as a multi-purpose medicinal agent, while *Leonotis nepetifolia* has been used successfully in Siddha medicine for centuries, more clinical trials should be conducted to support its therapeutic use. Moreover, the therapeutic potential of the plant should also be checked when used in combination with other herbal drugs.^{1, 5, 8, 9, 13}.

CONCLUSION

Ethno pharmacological activity and traditional uses of natural compounds, especially of plant origin received much attention in recent years as they are well tested for their efficacy and general believed to be safe for human use. Traditionally, plants are used in the treatment of many infections and systemic disorders. More than hundreds of chemical compounds are derived from plants which have medicinal values due to their health-enhancing and therapeutic properties are referred as herbs. Through screening of literature available on *Leonotis nepetifolia* depicted the fact that it is a popular remedy among the various ethnic groups Siddha and Ayurvedic properties.

REFERENCE

- Darley Maria Oliveira, Fernanda Germano Melo, Sikiru OlaitanBalogun, Adriana Flach, Edineide Cristina Alexandre de Souza, Gilmar Prado de Souza, Iolanda do NascimentoAraujo Rocha, Luiz Antonio MendonçaAlvesda Costa, Ilsamar Mendes Soares, Larissa Irene da Silva, Sergio DonizetiAscencio andDomingosTabajara de Oliveira Martins.Antibacterial mode of action of the hydroethanolic extract of*Leonotisnepetifolia*(L.) R. Br. involves bacterial membrane perturbations, *Journal of Ethnopharmacology*, 2015; 17 (2): 356-363.
- Ndukui James Gakunga, Godfrey Kateregga, Larry Fred Sembajwe and John Kateregga. Antidiarrheal activity and phytochemical profile of the ethanolic leaf extract of *Leonotisnepetifolia*(Lion's ear) in Wistar albino rats. *Journal of Intercultural Ethnopharmacology*,2013; 2(2):121-126.
- Nidhi Gauba Dhawan, Ambrina Sardar Khan and Prateek Srivastava. A General Appraisal of *Leonotisnepetifolia*(L) R. Br. An Essential Medicinal Plant, Bulletin of Environment. *Pharmacology and Life Sciences*, 2013; 2 (8): 118-121.
- 4. Maliheh Najari Beidokhti and PrakashHS. Antioxidant and Anti-inflammatory potential of selected medicinal plantof*Lamiaceae* family. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2013; 9(5): 85-89.
- Tiziana Fornari, Alejandro Ruiz-Rodriguez, Gonzalo Vicente, Erika Vazquez, Monica R, Garcia-Risco and Guillermo Reglero. Kinetic study of the supercritical co-extraction of different plants from *Lamiaceae*family.*Journal of Supercritical Fluids*, 2012; 6 (4): 1-8.
- 6. Syed Imran, Suradkar SS and Koche DK. Phytochemical analysis of *leonotisnepetifolia*(l.)
 R. BR.a wild medicinal plant of *Lamiaceae*, *Bioscience discovery*, 2012;3(2): 197-196
- Ana Paula de Oliveira, Amanda LeiteGuimaraes, EdigeniaCavalcante da Cruz Araujo1, Izabel Cristina Casanova Turatti, Norberto Peporine Lopes and Jackson Roberto Guedes da

Silva Almeida. GC-MS analysis of esterified fatty acids obtained from leaves of wild and cultivated specimens *Leonotisnepetifolia.Journal of Medicinal Plants Research*, 2015; 9 (16): 25-34.

- Sridharan Gurunagarajan and Brindha Pemaiah. Comparative studies on cytotoxic effect of Hyptissuaveolens Poit and Leonotisnepeatefolia R.Br. against EAC cell lines. Journal of Pharmacy Research, 2011; 4 (4): 1222-1224.
- 9. Usharani Veerabadran, Anuradha Venkatraman, Aroumougame Souprayane, Mathivanan Narayanasamy, Dhanalakshmi Perumal, Sagadevan Elumalai, Sindhu Sivalingam, Vadivelu Devaraj and Arumugam Perumal. Evaluation of antioxidant potential of leaves of *Leonotisnepetifolia* and its inhibitory effect on MCF7 and Hep2 cancer cell lines. *Asian Pacific Journal of Tropical Disease*, 2013; 7(3):103-110.
- 10. Williams AF, Clement YN, Nayak SB and RaoAVC.*Leonotisnepetifolia* protects against acetaminophen-induce hepatotoxicity: Histological Studies and the role of antioxidant enzymes,*Natural Products Chemistry & Research*, 2016; 9 (4):32-35.
- 11. Reshmi Pushpan, Reshmi Pushpan, Mukesh B, Nariya and Ashok BK. Evaluation of antiarthritic potential of *Leonotis nepetifolia*R.Br. againstfreund's adjuvant induced arthritis. *Journal of ayurvedaand integrated medical sciences*, 2017; 7 (2): 89-94.
- Danuta Sobolewska, Paweł Pasko, AgnieszkaGalanty, Justyna Makowska-Wąs, KingaPadło and Wojciech Wasilak. Preliminary phytochemical and biological screening of methanolic and acetone extracts from *Leonotisnepetifolia*(L.) R. Br. *Journal of Medicinal Plants Research*, 2012;6(30):4582-4585.
- 13. Sarath Chandirian, Musale JayeshVasudeo, Chaudharia PD. Evaluation of spermato toxicity of *Leonotisnepetifolia* in male rats.*International journal of research in pharmaceutical sciences*, 2017; 8(4): 585-589.
- 14. Subir Kumar Khawas and MishraPK.Ecology of *Leonotisnepetifolia* L. in Lodna Area of jharia coal field.*International journal of bioassays*, 2015; 4(10):4448-4450.
- 15. Udaya Prakash, Bhuvaneswari, Balamurugan A, VaishnaviS, sunishasugunan, sindhu Meena, Chandran M and Sekar H. Studies on antibacterial, larvicidal, pesticidal activities and phytochemistry of *Leonotisnepetifolia* (Linn) R.Br. *IJRPS*, 2013;4(2): 303-309.

- 16. Trivedi Ashish, Sethiya Neeraj K and Mishra SH. Preliminary pharmacognostic and phytochemical analysis of Grantbika (Leonotisnepetifolia): An Ayurvedic herb *.Indian journal of traditional knowledge*, 2011;10(4):682-688
- 17. Aye Aye Maw. Structural elucidation of an unknown bioactive organic compound isolated from Myanmar traditional indigenous medicinal plant *Leonotisnepetifolia* (L) R.Br (ame da zoe pin). *Universities research journal*, 2011; 4(3):87-94.
- 18. Gupta R and Gupta GD. Determination of plant constituents in fractions of *cordiaobliqua*willd. Leaf methanolic extract using GC-MS analysis. *International journal of pharmacognosy and phytochemical research*, 2017; 9(9): 1274-1279.
- 19. AmuthaIswarya Devi J and Kottaimuthu A, Gas chromatography-mass spectrometry analysis of bioactive constituents in the ethanolic extract of *Saccharumspontaneum*linn. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2014; 6(2): 975-983.
- 20. Aswal BS, Goel AK, Kulshrestha DK, Mehrotra BN and Patnaik GK. Screening of Indian plants for biological activity: Part XV. *Indian J Exp Biol.* 1996; 3(4): 444–67.
- 21. Prakash Ramakrishnan, Dhivya Ramadoss, Priyadarshini Muthulingam, Ramya Nedunchezhian and Karthick Krishna moorthy. Anti-diabetic, Anti-hyperlipidemic, Antioxidant Property of *Cordiaobliqua* on Streptozotocin Induced Diabetic Rats, *J Young Pharm.* 2017; 9(3): 321-326.
- 22. Ameera Omran Hussein, Ghaidaa Jihadi Mohammed, Mohammed Yahya Hadi and Imad Hadi Hameed. Phytochemical screening of methanolic dried galls extract of *Quercusinfectoria* using gas chromatography-mass spectrometry (GC-MS) and Fourier transform-infrared (FT-IR), *Journal of Pharmacognosy and Phytotherapy*. 2016; 8(3): 49-59.
- 23. Mohanad Jawad Kadhim, Azhar Abdulameer Sosa and Imad Hadi Hameed. Evaluation of anti-bacterial activity and bioactive chemical analysis of *Ocimumbasilicum*using Fourier transform infrared (FT-IR) and gas chromatography-mass spectrometry (GC-MS) techniques, *Journal of Pharmacognosy and Phytotherapy*. 2016; 8(6): 127-146.
- 24. Sermakkani M and Thangapandian V. GC-MS Analysis of cassia italica leaf methanol extract, *Asian Journal of Pharmaceutical and Clinical Research*. 2012; 5(2): 562-570.

- 25. Bagavathi Perumal Ezhilanand Ramasamy Neelamegam. GC-MS analysis of phytocomponents in the ethanol extract of *PolygonumchinenseL*, *Pharmacognosy Res*. 2012; 4(1): 11–14.
- 26. Gopalakrishnan S. GC-MS analysis of some bioactive constituents of *Mussaenda frondosa* linn, *International Journal of Pharma and Bio Sciences*. 2011; 2(1): 155–163.
- 27. Hema R, Kumaravel S and Alagusundaram K.GC/MS determination of bioactive components of *Murrayakoenigii, Journal of American Science*. 2011; 7(1): 345-352.
- 28. Jeyaraman Amutha Iswarya Devi, Velayutham Mani Mala and Narayanan Venkateshan. Review on Cordia obliqueWilld. International Journal of Chemical and Pharmaceutical Sciences. 2019;10(2): 26-30.